INDEX

TO VOLUME 35

SUBJECT INDEX

Abortion
A blow for abortion law appeal, FYI, p. 60.

Abortion incidence and services in the United States in 2000, Finer LB and Henshaw SK, p. 6.

Abortion restrictions in Russia grow, FYI, p. 201.

Abortion: teaching why as well as how, Viewpoint, Stewart FH and Darney PD, p. 37.

Accessibility of abortion services in the United States, 2001, Henshaw SK and Finer LB, p. 16.

Another early abortion option, FYI, p. 5.

Answering the call, FYI, p. 160.

Beyond apocalypse and apology: a moral defense of abortion, Viewpoint, Borgmann C and Weiss C, p. 40.

Complication rate is lower for surgical than medical second-trimester abortion, Digest, p. 50.

Convincing new providers to offer medical abortion: what will it take? Viewpoint, Coeytaux F, Moore K and Gelberg L, p. 44.

Foul! FYI, p. 112.

Public health impact of legal abortion: 30 years later, Comment, Cates W, Jr., Grimes DA and Schulz KF, p. 25.

Reflections of a provider before and since *Roe*: from the Voices of Choice Archive, Viewpoint, Physicians for Reproductive Choice and Health, p. 34.

Roe v. Wade at 30: what are the prospects for abortion provision? Viewpoint, Joffe C, p. 29.

Shortly after early abortion, women report significant improvements in physical and emotional well-being, Digest, p. 150.

Barrier Methods
A new look at an old method: the diaphragm, Viewpoint, Harvey SM, Bird ST and Branch MR, p. 270.

Evaluation of the efficacy of a nonlatex condom: results from a randomized, controlled clinical trial, Walsh TL, Frezieres RG, Peacock K, Nelson AL, Clark VA and Bernstein L, p. 79.

Family planning providers' perspectives on dual protection, see **Service Providers.**

High proportions of college men using condoms report errors and problems, Digest, p. 50.

How do you use this thing? FYI, p. 200.

Method-related problems account for most failures of the female condom, Digest, p. 193.

Sponge bounces back, FYI, p. 61.

Breast-Feeding
Breast-feeding is catching on, FYI, p. 200.

Clinician support may play a role in mothers' decision to continue breast-feeding, Digest, p. 278.

Contraceptive Practice
A new look at an old method: the diaphragm, see **Barrier Methods.**

Choosing does not mean using, FYI, p. 244.

High proportions of college men using condoms report errors and problems, see **Barrier Methods.**

Partner influence on early discontinuation of the pill in a predominantly Hispanic population, Kerns J, Westhoff C, Morroni C and Murphy PA, p. 256.

Patterns of contraceptive use within teenagers' first sexual relationships, see **Teenagers.**

Contraceptive Research
Evaluation of the efficacy of a nonlatex condom: results from a randomized, controlled clinical trial, see **Barrier Methods.**

Inching toward a male method, FYI, p. 245.

Emergency Contraception
Availability of emergency contraception through student health centers is growing, but gaps remain, Digest, p. 54.

Catholic hospitals offer limited access to emergency pills, FYI, p. 5.

Emergency contraception use increases when pills are provided in advance, Digest, p. 279.

Three differing emergency contraceptive regimens are equally effective, Digest, p. 152.

Variations on a theme by Yuzpe, FYI, p. 160.

Evaluation of Programs
An evaluation of California's Adolescent Sibling Pregnancy Prevention Program, East P, Kiernan E and Chávez G, p. 62.

Government Policy, Programming and Funding
Government for the people, FYI, p. 4.

Need information? don't ask the government, FYI, p. 5.

Newspeak 2003, FYI, p. 112.

Hormone Therapy
Study results have left many women confused about hormone therapy, Digest, p. 149.

International
British invasion, FYI, p. 161.

Cairo agreement stands, despite U.S. opposition, FYI, p. 4.

High rates of congenital syphilis in Russia linked to inadequate prenatal care, see **Sexually Transmitted Diseases.**

How to make a quick euro, FYI, p. 160.

UK youth prefer peer-led sexuality education classes to teacher-led programs, see **Teenagers.**

When and why do young people in the United Kingdom first use sexual health services? see **Teenagers.**

Intimate Partner Violence
History repeats itself, FYI, p. 245.

Women have different risk factors for verbal, physical partner abuse, Digest, p. 106.

Intrauterine Devices
IUD-related knowledge, attitudes and practices among Navajo Area Indian Health Service providers, Espey E, Ogburn T, Espey D and Etsitty V, p. 169.

Males
Heterosexually active men's beliefs about methods for preventing sexually transmitted diseases, Gillmore MR, Stielstra S, Huang B, Baker SA, Beadnell B and Morrison DM, p. 121.

Integrating *Chlamydia trachomatis* control services for males in female reproductive health programs, Viewpoint, McConnell KJ, Packel L, Biggs MA, Chow JM and Brindis C, p. 226.

Making males mindful of their sexual and procreative identities: using self-narratives in field settings, Viewpoint, Marsiglio W, p. 229.

Man2Man: a promising approach to addressing the sexual and reproductive health needs of young men, Special Report, Sherrow G, Ruby T, Braverman PK, Bartle N, Gibson S and Hock-Long L, p. 215.

Where does reproductive health fit into the lives of adolescent males? Marcell AV, Raine T and Eyre SL, p. 180.

Services for men at publicly funded family planning agencies, 1998–1999, Finer LB, Darroch JE and Frost JJ, p. 202.

The other half of the equation: serving young men in a young women's reproductive health clinic, Raine T, Marcell AV, Rocca CH and Harper CC, p. 208.

The Young Men's Clinic: addressing men's reproductive health rights and responsibilities, Special Report, Armstrong B, p. 220.

Marriage and the Family
Having two parents helps, FYI, p. 60.

Maternal and Child Health
Effects of psychosocial risk factors and prenatal interventions on birth weight: evidence from New Jersey's HealthStart Program, Reichman NE and Teitler JO, p. 130.

How's the baby? FYI, p. 113.

Maternal mortality exceeds U.S. goal; age and racial differences are marked, Digest, p. 189.

Mothers of many newborns with HIV missed chances to obtain preventive care, Digest, p. 188.

Neighborhood-level effects on babies' birth weight differ by race in Chicago, Digest, p. 153.

Youngest mothers' infants have greatly elevated risk of dying by age one, Digest, p. 52.

New Methods
Contraceptive ring found safe, effective; most users would recommend it, Digest, p. 53.

Oral Contraceptives
Partner influence on early discontinuation of the pill in a predominantly Hispanic population, see **Contraceptive Practice.**

Pill use protects against ovarian cancer; hormone therapy increases risk, Digest, p. 56.

Relative risk of cervical cancer rises with duration of oral contraceptive use, Digest, p. 192.

Some women with genetic susceptibility to breast cancer face elevated risk from oral contraceptive use, Digest, p. 105.

Women's pill IQ is lacking, FYI, p. 201.

Population Policy and Research
Measurement and meaning of unintended pregnancy, Comment, Santelli J, Rochat R, Hatfield-Timajchy K, Gilbert BC, Curtis K, Cabral R, Hirsch JS, Schieve L and other members of the Unintended Pregnancy Working Group, p. 94.

Pregnancy and Childbirth
Boys take more energy, FYI, p. 200.

Despite profile suggesting low risk, Indian-born U.S. mothers have high levels of some poor birth outcomes, Digest, p. 276.

Early prenatal care may not ensure improved outcomes among poor, rural infants, Digest, p. 275.

Effects of psychosocial risk factors and prenatal interventions on birth weight: evidence from New Jersey's HealthStart Program, see **Maternal and Child Health.**

For low-risk women, care from a nurse-midwife is a safe option in pregnancy, Digest, p. 275.

For whites, but not blacks, binge drinking is linked to unintended pregnancy, Digest, p. 236.

High rates of congenital syphilis in Russia linked to inadequate prenatal care, see **Sexually Transmitted Diseases.**

Not all Ohio physicians offer HIV testing during standard prenatal care, Digest, p. 194.

Odds of ectopic pregnancy are sharply elevated among heavy smokers, Digest, p. 151.

Partner violence during pregnancy increases risk of adverse outcomes, Digest, p. 238.

Racial disparities in early and adequate prenatal care decreased among U.S. women in 1980s and 1990s, Digest, p. 103.

Too many cesareans? FYI, p. 201.

What would Ford think? FYI, p. 161.

Wine, women and pregnancy, FYI, p. 244.

Public Opinion
Wouldn't it be nice? FYI, p. 161.

Religion
Religious teenagers may have a lowered risk of engaging in unsafe sexual behavior, see **Teenagers.**

Teenage women who are devoted to their religion have reduced sexual risk, see **Teenagers.**

Vatican explains it all, FYI, p. 60.

Reproductive Health
Black women and menopause, FYI, p. 61.

Cervical cancer risk rises if women with HPV also have herpes infection, Digest, p. 108.

Hysterectomy rate among Hispanic women is only half that of white women, Digest, p. 155.

More harm than good, FYI, p. 113.

New cervical cancer screening guidelines, FYI, p. 5.

Pill use protects against ovarian cancer; hormone therapy increases risk, see **Oral Contraceptives.**

Relative risk of cervical cancer rises with duration of oral contraceptive use, see **Oral Contraceptives.**

Some women with genetic susceptibility to breast cancer face elevated risk from oral contraceptive use, see **Oral Contraceptives.**

Study results have left many women confused about hormone therapy, see **Hormone Therapy.**

Young women and Pap smears, FYI, p. 113.

Young women gain on cancer, FYI, p. 245.

Service Providers
Access to adolescent reproductive health services: financial and structural barriers to care, see **Teenagers.**

Accessibility of abortion services in the United States, 2001, see **Abortion.**

A new look at an old method: the diaphragm, see **Barrier Methods.**

Clinician support may play a role in mothers' decision to continue breast-feeding, see **Breast-Feeding.**

Convincing new providers to offer medical abortion: what will it take? see **Abortion.**

Family planning providers' perspectives on dual protection, Mantell JE, Hoffman S, Exner TM, Stein ZA and Atkins K, p. 71.

Few U.S. school-based health centers offer contraceptives on-site, Digest, p. 239.

Integrating *Chlamydia trachomatis* control services for males in female reproductive health programs, see **Males.**

IUD-related knowledge, attitudes and practices among Navajo Area Indian Health Service providers, see **Intrauterine Devices.**

Man2Man: a promising approach to addressing the sexual and reproductive health needs of young men, see **Males.**

Not all Washington-area primary care practices offer all needed adolescent reproductive health services, see **Teenagers.**

Practice, practice, practice, FYI, p. 161.

Reflections of a provider before and since *Roe:* from the Voices of Choice Archive, see **Abortion.**

Roe v. Wade at 30: what are the prospects for abortion provision? see **Abortion.**

Services for men at publicly funded family planning agencies, 1998–1999, see **Males.**

The other half of the equation: serving young men in a young women's reproductive health clinic, see **Males.**

The Young Men's Clinic: addressing men's reproductive health rights and responsibilities, see **Males.**

Sex Education
Factors associated with the content of sex education in U.S. public secondary schools, Landry DJ, Darroch JE, Singh S and Higgins J, p. 261.

Teenagers with mental disability lack reproductive education and knowledge; still, many have had sex, see **Teenagers.**

UK youth prefer peer-led sexuality education classes to teacher-led programs, see **Teenagers.**

Sexual Behavior
Adolescent drinking and sex: findings from a daily diary study, see **Teenagers.**

Among female adolescents, STD history is associated with demographically or socially dissimilar partnering, see **Teenagers.**

Even after having an STD, many teenagers do not adopt safer-sex practices, see **Teenagers.**

Family connectedness and sexual risk-taking among urban youth attending alternative high schools, see **Teenagers.**

Improving women's sexual assertiveness, see **Teenagers.**

Large proportions of men and women with HIV have sex without telling partners that they are infected, Digest, p. 235.

Parents' smoking, seat belt nonuse may be linked to increased odds of adolescents' sexual debut, see **Teenagers.**

Preventing sexual risk behaviors and pregnancy among teenagers: linking research and programs, see **Teenagers.**

Reducing risk among men who have sex with men, FYI, p. 161.

Religious teenagers may have a lowered risk of engaging in unsafe sexual behavior, see **Teenagers.**

Sex in the countryside, FYI, p. 61.

Sexual behavior is safer when students can get condoms at their schools, see **Teenagers.**

Teenage women who are devoted to their religion have reduced sexual risk, see **Teenagers.**

Teenagers have the same risks whether partners are sequential or concurrent, see **Teenagers.**

Teenagers with the least adult supervision engage in the most sexual activity, see **Teenagers.**

Timing, amount of teenage alcohol or marijuana use may make future risky sex more likely, Digest, p. 49.

Youth who first got drunk by age 12 have relative-

ly high levels of risky sexual behavior during college, see **Teenagers.**

Sexually Transmitted Diseases

A new tack on HIV prevention efforts, FYI, p. 113.

About half of U.S. adults have had an HIV test, but testing often occurs late in the course of infection, Digest, p. 274.

Acquiring a new partner is linked to increased HPV risk among young women, Digest, p. 150.

Cervical cancer risk rises if women with HPV also have herpes infection, see **Reproductive Health**.

Disease burden for blacks, FYI, p. 201.

Disease risk is elevated among people who feel bad about having sex, Digest, p. 239.

Emergency-room visits can be good opportunities for sexually transmitted disease screening, Digest, 191.

Foreign-born STD clinic clients with HIV likely acquired infection here, Digest, p. 104.

Heterosexually active men's beliefs about methods for preventing sexually transmitted diseases, see **Males**.

High rates of congenital syphilis in Russia linked to inadequate prenatal care, Digest, p. 190.

Large proportions of men and women with HIV have sex without telling partners that they are infected, see **Sexual Behavior.**

Learning who stops care, FYI, p. 201.

Mothers of many newborns with HIV missed chances to obtain preventive care, see **Maternal and Child Health**.

Not all Ohio physicians offer HIV testing during standard prenatal care, see **Pregnancy and Childbirth**.

Reinfections make up a large share of gonorrhea cases diagnosed at clinics, Digest, p. 237.

Screening opportunities missed, FYI, p. 244.

Survival means new risks, FYI, p. 112.

Treating common vaginal infections may lower women's herpes risk, Digest, p. 277.

What the British don't know could hurt them, FYI, p. 4.

You think you know a person, FYI, p. 245.

Sterilization

Child disability and mothers' tubal sterilization, Park JM, Hogan DP and Goldscheider FK, p. 138.

Hysterectomy rate among Hispanic women is only half that of white women, Digest, p. 155.

Teenagers

Access to adolescent reproductive health services: financial and structural barriers to care, Viewpoint, Hock-Long L, Herceg-Baron R, Cassidy AM and Whittaker PG, p. 144.

Adolescent drinking and sex: findings from a daily diary study, Morrison DM, Gillmore MR, Hoppe MJ, Gaylord J, Leigh BC and Rainey D, p. 162.

Among female adolescents, STD history is associated with demographically or socially dissimilar partnering, Digest, p. 234.

An evaluation of California's Adolescent Sibling Pregnancy Prevention Program, see **Evaluation of Programs**.

Circle of life, FYI, p. 61.

Despite having received relevant education, youth lack knowledge of STDs, Digest, p. 51.

Dual use grows among teenagers, FYI, p. 244.

Even after having an STD, many teenagers do not adopt safer-sex practices, Digest, p. 103.

Factors associated with the content of sex education in U.S. public secondary schools, see **Sex Education**.

Family connectedness and sexual risk-taking among urban youth attending alternative high schools, Markham CM, Tortolero SR, Escobar-Chaves SL, Parcel GS, Harrist R and Addy RC, p. 174.

Going to the chapel, FYI, p. 5.

HIV programs for teenagers work, FYI, p. 160.

How to win friends...? FYI, p. 200.

How well do pharmacies serve teenagers? FYI, p. 112.

Improving women's sexual assertiveness, Letter, Tolman DL, p. 48.

Is sex depressing? FYI, p. 4.

Lessons not learned, FYI, p. 60.

Mothers exert more influence on timing of first intercourse among daughters than among sons, Digest, p. 55.

Not all Washington-area primary care practices offer all needed adolescent reproductive health services, Digest, p. 148.

Parents' smoking, seat belt nonuse may be linked to increased odds of adolescents' sexual debut, Digest, p. 102.

Patterns of contraceptive use within teenagers' first sexual relationships, Manlove J, Ryan S and Franzetta K, p. 246.

Preventing sexual risk behaviors and pregnancy among teenagers: linking research and programs, Comment, Kalmuss D, Davidson A, Cohall A, Laraque D and Cassell C, p. 87.

Providers do not fully use adolescent well-care visits to discuss sexual health, Digest, p. 188.

Rap-related risk, FYI, p. 113.

Religious teenagers may have a lowered risk of engaging in unsafe sexual behavior, Digest, p. 279.

Sexual behavior is safer when students can get condoms at their schools, Digest, p. 236.

Teenage women who are devoted to their religion have reduced sexual risk, Digest, p. 107.

Teenagers have the same risks whether partners are sequential or concurrent, Digest, p. 192.

Teenagers with mental disability lack reproductive education and knowledge; still, many have had sex, Digest, p. 187.

Teenagers with the least adult supervision engage in the most sexual activity, Digest, p. 106.

Timing, amount of teenage alcohol or marijuana use may make future risky sex more likely, see **Sexual Behavior**.

UK youth prefer peer-led sexuality education classes to teacher-led programs, Digest, p. 109.

When and why do young people in the United Kingdom first use sexual health services? Stone N and Ingham R, p. 114.

Where does reproductive health fit into the lives of adolescent males? see **Males**.

Youth who first got drunk by age 12 have relatively high levels of risky sexual behavior during college, Digest, p. 154.

AUTHOR INDEX

Addy RC, see Markham CM.

Armstrong B, The Young Men's Clinic: addressing men's reproductive health rights and responsibilities, Special Report, p. 220.

Atkins K, see Mantell JE.

Baker SA, see Gillmore MR.

Bartle N, see Sherrow G.

Beadnell B, see Gillmore MR.

Bernstein L, see Walsh TL.

Biggs MA, see McConnell KJ.

Bird ST, see Harvey SM.

Borgmann C and Weiss C, Beyond apocalypse and apology: a moral defense of abortion, Viewpoint, p. 40.

Branch MR, see Harvey SM.

Braverman PK, see Sherrow G.

Brindis C, see McConnell KJ.

Cabral R, see Santelli J.

Cassell C, see Kalmuss D.

Cassidy AM, see Hock-Long L.

Cates W, Jr., Grimes DA and Schulz KF, Public health impact of legal abortion: 30 years later, Comment, p. 25.

Chávez G, see East P.

Chow JM, see McConnell KJ.

Clark VA, see Walsh TL.

Coeytaux F, Moore K and Gelberg L, Convincing new providers to offer medical abortion: what will it take? Viewpoint, p. 44.

Cohali A, see Kalmuss D.

Curtis K, see Santelli J.

Darney PD, see Stewart FH.

Darroch JE, see Finer LB; Landry DJ.

Davidson A, see Kalmuss D.

East P, Kiernan E and Chávez G, An evaluation of California's Adolescent Sibling Pregnancy Prevention Program, p. 62.

Escobar-Chaves SL, see Markham CM.

Espey D, see Espey E.

Espey E, Ogburn T, Espey D and Etsitty V, IUD-related knowledge, attitudes and practices among Navajo Area Indian Health Service providers, p. 169.

Etsitty V, see Espey E.

Exner TM, see Mantell JE.

Eyre SL, see Marcell AV.

Finer LB and Henshaw SK, Abortion incidence and services in the United States in 2000, p. 6.

Finer LB, Darroch JE and Frost JJ, Services for men at publicly funded family planning agencies, 1998–1999, p. 202.

Finer LB, see Henshaw SK.

Franzetta K, see Manlove J.

Frezieres RG, see Walsh TL.

Frost JJ, see Finer LB.

Gaylord J, see Morrison DM.

Gelberg L, see Coeytaux F.

Gibson S, see Sherrow G.

Gilbert BC, see Santelli J.

Gillmore MR, Stielstra S, Huang B, Baker SA, Beadnell B and Morrison DM, Heterosexually active men's beliefs about methods for preventing sexually transmitted diseases, p. 121.

Gillmore MR, see Morrison DM.

Goldscheider FK, see Park JM.

Grimes DA, see Cates W, Jr.

Harper CC, see Raine T.

Harrist R, see Markham CM.

Harvey SM, Bird ST and Branch MR, A new look at an old method: the diaphragm, Viewpoint, p. 270.

Hatfield-Timajchy K, see Santelli J.

Henshaw SK and Finer LB, Accessibility of abortion services in the United States, 2001, p. 16.

Henshaw SK, see Finer LB.

Herceg-Baron R, see Hock-Long L.

Higgins J, see Landry DJ.

Hirsch JS, see Santelli J.

Hock-Long L, Herceg-Baron R, Cassidy AM and Whittaker PG, Access to adolescent reproductive health services: financial and structural barriers to care, Viewpoint, p. 144.

Hock-Long L, see Sherrow G.

Hoffman S, see Mantell JE.

Hogan DP, see Park JM.

Hoppe MJ, see Morrison DM.

Huang B, see Gillmore MR.

Ingham R, see Stone N.

Joffe C, *Roe v. Wade* at 30: what are the prospects for abortion provision? Viewpoint, p. 29.

Kalmuss D, Davidson A, Cohall A, Laraque D and Cassell C, Preventing sexual risk behaviors and pregnancy among teenagers: linking research and programs, Comment, p. 87.

Kerns J, Westhoff C, Morroni C and Murphy PA, Partner influence on early discontinuation of the pill in a predominantly Hispanic population, p. 256.

Kiernan E, see East P.

Landry DJ, Darroch JE, Singh S and Higgins J, Factors associated with the content of sex education in U.S. public secondary schools, p. 261.

Laraque D, see Kalmuss D.

Leigh BC, see Morrison DM.

Manlove J, Ryan S and Franzetta K, Patterns of contraceptive use within teenagers' first sexual relationships, p. 246.

Mantell JE, Hoffman S, Exner TM, Stein ZA and Atkins K, Family planning providers' perspectives on dual protection, p. 71.

Marcell AV, Raine T and Eyre SL, Where does reproductive health fit into the lives of adolescent males? p. 180.

Marcell AV, see Raine T.

Markham CM, Tortolero SR, Escobar-Chaves SL, Parcel GS, Harrist R and Addy RC, Family connectedness and sexual risk-taking among urban youth attending alternative high schools, p. 174.

Marsiglio W, Making males mindful of their sexual and procreative identities: using self-narratives in field settings, Viewpoint, p. 229.

McConnell KJ, Packel L, Biggs MA, Chow JM and Brindis C, Integrating *Chlamydia trachomatis* control services for males in female reproductive health programs, Viewpoint, p. 226.

Moore K, see Coeytaux F.

Morrison DM, Gillmore MR, Hoppe MJ, Gaylord J, Leigh BC and Rainey D, Adolescent drinking and sex: findings from a daily diary study, p. 162.

Morrison DM, see Gillmore MR.

Morroni C, see Kerns J.

Murphy PA, see Kerns J.

Nelson AL, see Walsh TL.

Ogburn T, see Espey E.

Packel L, see McConnell KJ.

Parcel GS, see Markham CM.

Park JM, Hogan DP and Goldscheider FK, Child disability and mothers' tubal sterilization, p. 138.

Peacock K, see Walsh TL.

Physicians for Reproductive Choice and Health, Reflections of a provider before and since *Roe*: from the Voices of Choice Archive, Viewpoint, p. 34.

Raine T, Marcell AV, Rocca CH and Harper CC, The other half of the equation: serving young men in a young women's reproductive health clinic, p. 208.

Raine T, see Marcell AV.

Rainey D, see Morrison DM.

Reichman NE and Teitler JO, Effects of psychosocial risk factors and prenatal interventions on birth weight: evidence from New Jersey's HealthStart Program, p. 130.

Rocca CH, see Raine T.

Rochat R, see Santelli J.

Ruby T, see Sherrow G.

Ryan S, see Manlove J.

Santelli J, Rochat R, Hatfield-Timajchy K, Gilbert BC, Curtis K, Cabral R, Hirsch JS, Schieve L and Other Members of the Unintended Pregnancy Working Group, The measurement and meaning of unintended pregnancy, Comment, p. 94.

Schieve L, see Santelli J.

Schulz KF, see Cates W, Jr.

Sherrow G, Ruby T, Braverman PK, Bartle N, Gibson S and Hock-Long L, Man2Man: a promising approach to addressing the sexual and reproductive health needs of young men, Special Report, p. 215.

Singh S, see Landry DJ.

Stein ZA, see Mantell JE.

Stewart FH and Darney PD, Abortion: teaching why as well as how, Viewpoint, p. 37.

Stielstra S, see Gillmore MR.

Stone N and Ingham R, When and why do young people in the United Kingdom first use sexual health services? p. 114.

Teitler JO, see Reichman NE.

Tolman DL, Improving women's sexual assertiveness, Letter, p. 48.

Tortolero SR, see Markham CM.

Unintended Pregnancy Working Group, see Santelli J.

Walsh TL, Frezieres RG, Peacock K, Nelson AL, Clark VA and Bernstein L, Evaluation of the efficacy of a nonlatex condom: results from a randomized, controlled clinical trial, p. 79.

Weiss C, see Borgmann C.

Westhoff C, see Kerns J.

Whittaker PG, see Hock-Long L.